

MASS SPECTRA OF RADIALLY AND ORBITALLY EXCITED STATES OF MESONS

V.Khruschov ¹, V.Savrin ², S.Semenov ³

^{a,c}RRC "Kurchatov Institute", Kurchatov Sq., 123182 Moscow, Russia

^bSkobeltsyn Institute of Nuclear Physics, Moscow State University

Abstract

Meson mass spectra, evaluated in the framework of the relativistic model of quasi-independent quarks, are presented. Mass values are obtained with the help of numerical calculations based on the Dirac equation and by phenomenological mass formulae. The Dirac equation involves the potential, which is sum of the vector quasi-Coulombic potential and the scalar linear rising confinement potential. The phenomenological mass formulae are applied to excited meson states consisting of u -, d -quarks and antiquarks with isotopical spin $I = 1$. A comparison of the evaluated mass spectra with existing data is performed. Problems of identification of some meson states in vector and scalar channels are discussed.

It is known that the calculation of hadron mass spectra on the level of experimental data precision [1] still remains among the unsolved QCD problems due to some technical and conceptual difficulties, which are related mainly with the nonperturbative effects, such as confinement and spontaneous chiral symmetry breaking. Nowadays hadron characteristics calculations are frequently carried out with the help of phenomenological models. Among phenomenological models the potential quark models are the simplest ones, they make it possible to represent the different stages of calculations in terms of commonly used physical quantities. Although these models are suitable rather well for the heavy quark hadrons treating of hadrons containing the light quarks is more complicated task, which demands relativistic and nonpotential effects to be taken into account [2, 3, 4]. The relativistic model of quasi-independent quarks is applicable to the description of properties of light and heavy hadrons [5, 6, 7, 8]. Below we present the results of calculations in the framework of this model [8, 9, 10] of mass spectra of quarkonium type mesons, which are orbitally and radially excited states. We compare results obtained with data, make predictions for mass values of unobserved yet meson states and discuss possible structures of some mesons in scalar and vector channels.

According to the main statement of the independent-quark model a hadron is considered as a system composed of a few non-interacting with each other directly valency constituents (quarks, diquarks and constituent gluons) with the coordinates $\mathbf{r}_i$, $i = 1, \dots, N$, which moving in some mean colour singlet confining field. It is assumed that the field is spherically symmetric and its motion in space is determined by motion of its center with the coordinate $\mathbf{r}_0$. We treat the mean field as a quasi-classical object possessing some energy ϵ_0 . Meantime each of N constituents interacting with the mean field gets the state with a definite value of its energy ϵ_i , which can be determined after solving of the model equation. We choose the Dirac equation as the model equation for quarks (the Klein-Gordon equation for diquarks and constituent gluons):

$$\sqrt{\lambda_i + m_i^2} \psi_i(\mathbf{r}_i) = [(\boldsymbol{\alpha}_i \mathbf{p}_i) + \beta_i(m_i + V_0) + V_1] \psi_i(\mathbf{r}_i), \quad (1)$$

with $E_i(n_i^r, j_i) = \sqrt{\lambda_i + m_i^2}$, $i = 1, 2$, $V_0(r) = \sigma r/2$ and $V_1(r) = -2\alpha_s/3r$, where the model parameters σ and α_s have meanings of the string tension and the strong coupling constant at small distances, correspondingly. In order to exclude superfluous meson states, the following selection rules for $n^{2S+1}L_J$ -states with quark masses m_1, m_2 and quantum numbers n_1^r, j_1, n_2^r, j_2 are used

$$\begin{aligned} n_1^r &= n_2^r = n - 1, & j_1 &= j_2 = J + 1/2, & \text{if } J &= L + S, \\ j_1 &= j_2 + 1 = J + 3/2, & \text{if } J &\neq L + S, & m_1 &\leq m_2, \end{aligned} \quad (2)$$

¹Talk given at 12th Lomonosov Conference on Elementary Particle Physics, Moscow, August 25-31, 2005

e-mail: khru@imp.kiae.ru

²e-mail: savrin@theory.sinp.msu.ru

³e-mail: semenov@imp.kiae.ru

After the elimination the angular dependence the radial equation obtained can be solved only by numerical methods. For calculating its eigenvalues the code algorithm, which was based on the Numerov three point recurrent relation [11], has been used jointly with the regularization procedure for the singular potential at $r = 0$. In order to estimate the spin-spin interaction between a quark and an antiquark we use the expression for this interaction from the relativized quark model [3]: $V_{S_1 S_2} = \frac{32\pi\alpha_s \mathbf{S}_1 \mathbf{S}_2}{9E_1 E_2} \delta^3(\mathbf{r})$, where $\mathbf{S}_1$ and $\mathbf{S}_2$ are the spin operators and E_1 , E_2 are the energies of quark and antiquark, respectively. Evaluation of the expectation value of this operator in the first order of perturbation theory shows that a contribution of the spin-spin interaction is a slowly varying function versus E_1 and E_2 and depends mainly on the total spin of quark-antiquark system and the α_s . For S-wave mesons we use the mass formula [9]:

$$M_m = E_{0m} + E_1(n_1^r, 1/2) + E_2(n_2^r, 1/2) + 4 \langle \mathbf{S}_1 \mathbf{S}_2 \rangle_{q_1 \bar{q}_2} V_{SS}, \quad (3)$$

where V_{SS} is approximately proportional to α_s with constant v_{0S} . As calculations show, the E_{0m} are negligible for the majority of 0^{-+} and 1^{--} mesons, excluding only a few cases, which will be discussed below. It should be noted, that for ground states of $\bar{q}q'$ -mesons the form of relation (3) is analogous (after the substitution $E_i \rightarrow m_i^c$, $i = 1, 2$) to the mass formula, which depends on constituent quark masses m_i^c [12]. Although for vector heavy mesons the contributions of spin-spin interaction are on the level of the model absolute uncertainty $\sim 40 MeV$, nevertheless, for pseudoscalar mesons these contributions should be taken into account. The magnitude of the spin-spin interaction constant v_{0S} is approximately equal to $100 MeV$. In the case of isoscalar mesons, which are neutral with respect to flavour quantum numbers, the terms due to the annihilation interaction for quark and antiquark or instanton interaction appear [13, 14]. But in the framework of our model the contributions of such type can be included into E_{0m} terms.

We have found that a suitable fit within the model accuracy for masses of ground state mesons, which are composed of quark and antiquark with u -, d -, s -, c - or b - flavours [1], is provided by the following values of E_{0m} parameter: $E_0(\eta_c) = E_c(J/\Psi) = -150 MeV$, $E_0(K) = -200 MeV$, $E_0(\pi) = E_0(\eta_b) = E_0(\Upsilon) = -450 MeV$. For the α_s and the quark masses we obtain the following values: $\bar{m} = (0.007 \pm 0.005)$ GeV, $\alpha_s^n = 0.60 \pm 0.15$, $m_s = (0.14 \pm 0.03)$ GeV, $\alpha_s^s = 0.47 \pm 0.10$, $m_c = (1.28 \pm 0.05)$ GeV, $\alpha_s^c = 0.33 \pm 0.03$, $m_b = (4.60 \pm 0.10)$ GeV, $\alpha_s^b = 0.27 \pm 0.02$, $\sigma = (0.20 \pm 0.01) GeV^2$. Within the model accuracy data for the well known experimental mass values for 1^{--} and 0^{-+} mesons with u -, d -, s -, c - or b quark and antiquark are consistent with the presented above model parameters. Thus in the framework of the model the flavor independence of the confinement potential [15, 16, 17] is confirmed as well as the concordance with the asymptotic freedom behaviour of the α_s . The evaluated mass spectra of mesons are displayed in the Table 1.

In spite of difficulties which exist in for evaluations of mass spectra of light hadrons a number of relations among their masses have been obtained using phenomenological considerations, such as the mass relations for Regge trajectories, daughter Regge trajectories [18, 19, 20, 21] and the mass formulae, obtained on the base of finite energy QCD sum rules [22]. In the framework of relativistic quasi-independent quarks model the quark and antiquark energies are determined from the solutions of the Dirac equation. However, for the superlight $\bar{q}q'$ -mesons with u - and d - quarks and antiquarks only, the phenomenological energy spectral functions $E_i(n_i, j_i, c, \kappa)$ are suitable within the relative uncertainty of the order of a few percents [7]:

$$M(n^{2S+1} L_J) = E_1(n_1^r, j_1, c, \kappa) + E_2(n_2^r, j_2, c, \kappa), \quad (4)$$

where

$$E_i(n_i, j_i, c, \kappa) = \begin{cases} c + \kappa \sqrt{2n^r + L + j - 1/2}, & L + j - 1/2 = 2k, \quad k = 0, 1, \dots \\ \kappa \sqrt{2n^r + L + j - 1/2}, & L + j - 1/2 = 2k + 1, \quad k = 0, 1, \dots \end{cases}$$

Note that nonpotential corrections, which are connected mainly with c value, are the most important for mass spectra of light mesons [13, 4]. When evaluating masses of unobserved yet excited meson states we use the formulae written above together with the values of two parameters c and

κ , which have been obtained by fitting the mass values of experimentally detected meson states. The c and κ values obtained by this manner are $c = 69 \text{ MeV}$, $\kappa = 382 \pm 4 \text{ MeV}$ [10]. The results obtained with $c = 69 \text{ MeV}$, $\kappa = 385 \text{ MeV}$ are shown in the Table 2, where the mass values of orbital excitations of superlight meson states up to $L = 4$ are presented.

The mass formulae presented above permit to explain the degeneracy of mass values for different J^{PC} mesons. For example, we get $M(\rho'') = M(\rho_3)$. If one substitute the data, then $1720 \pm 20 \text{ MeV} = 1688.8 \pm 2.1 \text{ MeV}$. In the high mass value region the degeneracy of such kind will be increased and it is seen in the $\sim 2315 \text{ MeV}$ region. As it follows from the Table 2, the mass value calculated for orbital excitation of the 1^{--} meson is equal to 1506 MeV . So mixing between the radial and orbital excitations for the 1^{--} mesons should take place, which leads to the experimentally visible vector meson states V'_R and V'_O . Taking into account the experimental data the following cases are most preferable. In the first case the observed bump at 1465 MeV consists of one resonance V'_R , that is mainly the radial excitation of ρ -meson. In the second case the observed bump at 1465 MeV consist of two resonances V'_R and V'_O (see also [23]). Another important problem is the interpretation of mesons discovered in scalar channel. The results of our evaluations (Table 2) confirm the existence of P -wave $\bar{q}q'$ -mesons with masses $\sim 980 \text{ MeV}$. Another fact in favour of P -wave $\bar{q}q'$ -mesons with masses $\sim 980 \text{ MeV}$ is the coincidence of the evaluated mass for the first radial excitation of $a_0(980)$ meson with the mass value of the $a_0(1450)$ meson [1]. This meson has been predicted for the first time in the framework of finite energy QCD sum rules [24].

References

- [1] S.Eidelman et al. (Particle Data Group), *Phys.Lett.* **B** 592, 1 (2004).
- [2] A.A.Bykov, I.M.Dremin, A.V.Leonidov, *Usp.Fis.Nauk* 143, 3 (1984).
- [3] S.Godfrey and N.Isgur, *Phys.Rev.* **D** 32, 189 (1985).
- [4] B.Diekmann, *Phys.Rep.* 159, 99 (1988).
- [5] P.N.Bogoliubov, *Ann.Inst. H.Poincare* 8, 2 (1968).
- [6] P.N.Bogoliubov, A.E.Dorokhov, *Part.& Nucl.* 8, 917 (1987).
- [7] V.V.Khrushchev, *Yad.Fiz.* 55, 773 (1992).
- [8] V.V.Khrushchev, V.I.Savrin, S.V.Semenov, *Phys.Lett.* **B** 525, 283(2002).
- [9] V.V.Khrushchev, S.V.Semenov, *Part.Nucl.Lett.* 5, 5 (2002).
- [10] V.V.Khrushchov, *prep. IAE-6359/2*, Moscow, 2005; [*hep-ph/0504077*].
- [11] B.V.Numerov, *Proc.Centr.Astrophys.Observ.* 2, 188 (1933).
- [12] Ya.B.Zeldovich, A.D.Sakharov, *Yad.Fiz.* 4, 395 (1966).
- [13] A.T.Filippov, *Yad. Fiz.* 29, 1035 (1979).
- [14] A.E.Dorokhov, Ya.A.Zubov, N.I.Kochelev, *Part.&Nucl.* 23 (1992) 1192.
- [15] A.E.Dorokhov, *prep. JINR R2-121159*, Dubna, 1979.
- [16] A.Martin, *prep. CERN TH-2741*, Geneva, 1979.
- [17] C.Quigg, J.L.Rosner, H.B.Thacker, *FERMILAB* 79-52(THY), 1979.
- [18] M.Toller in *Proc. 8 Nobel Symp. on Elem. Part. Theory*, N.Y. 1968.

- [19] G.Veneziano in Proc. Coral Gables Conf. on Fund. Inter., N.Y. 1969.
- [20] Y.Nambu in Lect. at the 1970 Copenhagen Summer Symp., 1970.
- [21] A.Anisovich, V.Anisovich, A.Sarantsev, *Phys.Rev.* **D** 62, 051502(2000).
- [22] A.L.Kataev, *arxiv: hep-ph/0406305*.
- [23] N.N.Achasov, A.A.Kozhevnikov, *Yad.Fiz.* 65, 158 (2002).
- [24] S.G.Gorishny, A.L.Kataev, S.A.Larin, *Phys.Lett.* **B** 135, 457 (1984).

Table 1.

Evaluated masses of the 1^{--} and 0^{-+} mesons in comparison with the data from Ref. [1].

Meson	$M_n^{exp}[MeV]$	$M_n^{th}[MeV]$	Meson	$M_n^{exp}[MeV]$	$M_n^{th}[MeV]$
ρ	775.8 ± 0.5	740	B	5279.2 ± 0.5	5250
ρ'	1465 ± 25	1455	B_s	5369.6 ± 2.4	5370
ρ''	1720 ± 20	1730	B_c	6400 ± 400	6450
ϕ	1019.456 ± 0.020	1010	η_b	-	9330
ϕ'	1680 ± 20	1650	π'	1300 ± 100	1290
ϕ''	-	2050	K'	-	1400
J/ψ	3096.916 ± 0.011	3060	D'	-	2450
ψ'	3686.093 ± 0.034	3650	D'_s	-	2560
ψ''	4040 ± 10	4070	η'_c	-	3600
ψ'''	4415 ± 6	4390	B'	-	5650
Υ	9460.30 ± 0.26	9470	B'_s	-	5750
Υ'	10023.26 ± 0.31	9990	B'_c	-	6800
Υ''	10355.2 ± 0.5	10325	η'_b	-	9960
Υ'''	10580 ± 3.5	10550	π''	1812 ± 14	1810
Υ^{5S}	10865 ± 8	10830	K''	-	1850
Υ^{6S}	11019 ± 8	10985	D''	-	2880
π	138.039 ± 0.004	120	D''_s	-	2950
K	495.66 ± 0.02	500	η''_c	-	3970
D	1867.0 ± 0.5	1850	B''	-	6070
D_s	1968.3 ± 0.5	1990	B''_s	-	6130
η_c	2979.6 ± 1.2	2990	B''_c	-	7150

Table 2.

Evaluated masses in MeV for the ground states and the orbital excitations of the $\bar{q}q'$ - mesons in comparison with the data from Ref.[1].

Meson	J^{PC}	$M^{exp}(MeV)$	$M^{ev}(MeV)$	Meson	J^{PC}	$M^{exp}(MeV)$	$M^{ev}(MeV)$
π	0^{-+}	138.039 ± 0.004	138 ± 30	ρ_3	3^{--}	1688.8 ± 2.1	1722 ± 30
ρ	1^{--}	775.8 ± 0.5	770 ± 30	a_2^3	2^{++}	-	1873 ± 30
a_0	0^{++}	984.7 ± 1.2	998 ± 30	b_3^3	3^{+-}	-	2024 ± 30
b_1	1^{+-}	1229.5 ± 3.2	1227 ± 30	a_3^3	3^{++}	-	2031 ± 30
a_1	1^{++}	1230 ± 40	1280 ± 30	a_4	4^{++}	2010 ± 12	2037 ± 30
a_2	2^{++}	1318.3 ± 0.6	1334 ± 30	a_3^4	3^{--}	-	2177 ± 30
a_1^2	1^{--}	-	1506 ± 30	b_4^4	4^{-+}	-	2316 ± 30
π_2	2^{-+}	1672.4 ± 3.2	1678 ± 30	a_4^4	4^{--}	-	2313 ± 30
a_2^2	2^{--}	-	1700 ± 30	a_5^4	5^{--}	-	2310 ± 30